

CONTROL ID: 1493571

TITLE: Testing seismic hazard models with Be-10 exposure ages for precariously balanced rocks

AUTHORS (FIRST NAME, LAST NAME): Dylan H. Rood^{1,2}, Rasool Anooshehpour³, Greg Balco⁴, Glenn P Biasi⁵, James N Brune⁵, Richard Brune⁶, Lisa Grant Ludwig⁶, Katherine J Kendrick⁷, Matthew Purvance^{8,5}, Inyo Saleeby²

INSTITUTIONS (ALL): 1. AMS Laboratory, Scottish Universities Environmental Research Centre (SUERC), East Kilbride, United Kingdom.

2. Earth Research Institute, University of California, Santa Barbara, CA, United States.

3. U.S. Nuclear Regulatory Commission, Washington, DC, United States.

4. Berkeley Geochronology Center, Berkeley, CA, United States.

5. Seismological Laboratory, University of Nevada, Reno, NV, United States.

6. Program in Public Health, University of California, Irvine, CA, United States.

7. Earthquake Science Center, US Geological Survey, Pasadena, CA, United States.

8. Itasca Consulting Group, Minneapolis, MN, United States.

ABSTRACT BODY: Currently, the only empirical tool available to test maximum earthquake ground motions spanning timescales of 10 ky-1 My is the use of fragile geologic features, including precariously balanced rocks (PBRs). The ages of PBRs together with their areal distribution and mechanical stability (“fragility”) constrain probabilistic seismic hazard analysis (PSHA) over long timescales; pertinent applications include the USGS National Seismic Hazard Maps (NSHM) and tests for ground motion models (e.g., Cybershake). Until recently, age constraints for PBRs were limited to varnish microlamination (VML) dating techniques and sparse cosmogenic nuclide data; however, VML methods yield minimum limiting ages for individual rock surfaces, and the interpretations of cosmogenic nuclide data were ambiguous because they did not account for the exhumation history of the PBRs or the complex shielding of cosmic rays. We have recently published a robust method for the exposure dating of PBRs combining Be-10 profiles, a numerical model, and a three-dimensional shape model for each PBR constructed using photogrammetry (Balco et al., 2011, Quaternary Geochronology). Here, we use our published method to calculate new exposure ages for PBRs at 6 sites in southern California near the San Andreas, San Jacinto, and Elsinore faults, including: Lovejoy Buttes (9 +/- 1 ka), Round Top (35 +/- 1 ka), Pacifico (19 +/- 1 ka, but with a poor fit to data), Beaumont South (17 +/- 2 ka), Perris (24 +/- 2 ka), and Benton Road (40 +/- 1 ka), in addition to the recently published age of 18.5 +/- 2.0 ka for a PBR at the Grass Valley site. We combine our ages and fragilities for each PBR, and use these data to test the USGS 2008 NSHM PGA with 2% in 50 year probability, USGS 2008 PSHA deaggregations, and basic hazard curves from USGS 2002 NSHM data.

KEYWORDS: [1150] GEOCHRONOLOGY / Cosmogenic-nuclide exposure dating, [7212] SEISMOLOGY / Earthquake ground motions and engineering seismology, [7221] SEISMOLOGY / Paleoseismology, [8111] TECTONOPHYSICS / Continental tectonics: strike-slip and transform.



Precariously balanced rock in southern California

(No Table Selected)

Additional Details

Previously Presented Material: 20% from Balco et al. (2011) in Quaternary Geochronology.

Contact Details

CONTACT (NAME ONLY): Dylan Rood

CONTACT (E-MAIL ONLY): Dylan.Rood@glasgow.ac.uk

TITLE OF TEAM:
